


GALAXY INTERACTIONS: OVERVIEW

François Schweizer
Carnegie Observatories, Pasadena

Introduction


- 1. Interactions in the Local Group**
- 2. Damaged Disks**
- 3. Ellipticals from Wrecked Disks**
- 4. I/A at High z and in Clusters**

Support: NSF (AST); NASA (StScI); Carnegie



Hubble 1936





- Why a dichotomy of shapes?
- Meaning of Hubble's morphological sequence?



Toomre 1977

- What dynamical processes drive galaxy evolution?
- Relative importance of interactions, minor mergers, & major mergers in shaping various types of galaxies?


Ellipticals from Wrecked Disks



NGC 3921, protoplanetary

Barnes (1999)

Simulated Merger of 2 Gas-Rich Disk Galaxies




Pericenter at $t=0$

Hibbard et al
(1999)

Signatures of Interactions

Morphological

Kinematic



Position-velocity
diagram

Optical & HI tails in NGC 4038/39

N 4038 / 39

N ↗

11"
 1 kpc

$D = 19 \text{ Mpc}$
 $H_0 = 75$




Whitmore
et al. (1999)

1.


INTERACTIONS AND MERGERS IN THE LOCAL GROUP

1. Evidence in Milky Way: Warp; retrograde orbits of stars and globular clusters; accretion of Sagittarius Dwarf; “spaghetti halo”?
2. The Magellanic Stream
3. M 31 and M 32
4. M 31 and us

Sagittarius Dwarf (Ibata, Gilmore, & Irwin 1994)



Velocities in Tidal Stream at 60° from Sgr I





Gardiner & Noguchi '96

Magellanic Stream: Comparison Simulations - Observations

Velocities

View on Sky

Magellanic Clouds and Stream simulation



M31 and Satellites (Sandage 61)



2.

DAMAGED DISKS

Central issues:

- How fragile are galaxy disks? (Tóth & Ostriker 92; Walker⁺96; Aguerri⁺01)
- Can bulges form through minor mergers? (Aguerre⁺01: “yes”)
- What fraction of bulges formed in this manner?

Current status:


1. Disks with gas are not as fragile as once thought
2. Minor mergers do occur in disk galaxies and seem to move them toward earlier Hubble types
3. Fraction of bulges built through mergers remains unknown; are there various sequences: disk first, bulge first, and/or mixed?

nature

125
YEARS
1869-1994

INTERNATIONAL WEEKLY JOURNAL OF SCIENCE

Volume 372 No. 6506 8 December 1994 \$8.50



Tracing galaxy interactions

Yun, Ho, & Lo (1994)


Polar Ring Galaxy NGC4650A



Rubin 94

NGC 4826 : Optical Velocities (Sat)


Slit position



Optical spectrum [NII]

H α

[NII]




- Velocity reversal
in gas is steep

- Stars corotate
with inner gas

- In transition reg,
evidence for gas
infall at ~ 100 km/s

LINE-OF-SIGHT VELOCITY (km/sec)



COUNTERROTATION IN DISK GALAXIES

Counterrotating Gas

S0: In 33 S0's: 14 co-, 3 skewed-, 9 counter-rotate;
⇒ >80% external origin (Bettoni⁺91; Bertola⁺99)

Sa: In 4 Sa's: Gas **decoupled** from stars (Haynes⁺00)

Sab: NGC 4826: **two** counterrotating gas systems
(Braun⁺92, 94; Rubin 94)

Counterrotating Disk Stars

S0: NGC 4550: stars rotate 50/50% pro/retrograde
(Rubin⁺92; Rix⁺92; Kuijken⁺96)


Sb: NGC 7217: stars rotate 70/30% pro/retrograde

Counter/Orthorotating Bulges!

Sa: NGC 4698: orthorotating bulge (Bertola⁺99)

Sb: NGC 7331, 2841: counterrotating bulges
(Prada⁺96, 99)

3.
ELLIPTICALS
FROM
WRECKED DISKS



SIMILARITIES: NGC 3921 & 7252 – ELLIPTICALS


- Single main bodies of $M_V \approx -22.0$ with $r^{1/4}$ -type light distributions ($\Rightarrow$ **violent relaxation**)
- **Power-law cores**, central lumin. density $>10^3 L_\odot/\text{pc}^3$
- Kinematically **decoupled core** (NGC 7252)
- *UBVI* color gradients typical of E's
- Velocity dispersions σ_V fit Faber-Jackson
- Fine structure (**shells**) from returning tidal material
- “**E+A**” **spectra** indicative of $b = 10\% - 30\%$ starbursts
- Globular-cluster populations increased by $>40\%$, $\sim 80\%$

$\Rightarrow$ N3921 & 7252 are 0.5 – 1.0 Gyr old protoellipticals.

TT72: **Merger rate** suggests that $\sim 1/3$ – all E's formed this way.

Hilbard & Mihos 95

Ripple ("Shell") Formation in DD Mergers



→ Prolonged return of tidal material forms loops and ripples for several Gyr.

Jedrzejewski & Schechter 88

Major-Axis Rotation Curves for 2 E's




FIG. 7. Stellar velocities as a function of nuclear distance in NGC 3608 (upper) and NGC 4494 (lower) from Jedrzejewski & Schechter (1988). Note that in NGC 3608 the discrete stellar core rotates retrograde; in NGC 4494 it rotates prograde.

(graph assembled by Rubin 94)

NGC 7252 (HST-4, Miller et al. 1997)



~500 candidate clusters $V \leq 26$ mag

32"
10 kpc

(HST)

From Giant Molecular Clouds to Globular Clusters Dichotomy of Shapes




M100 (Sc)

M87 (E0)

D. Malin - AAT

Ages of E+S0 Galaxies from H β (age sensitive) and Mg b, Fe




Results: - Field E have mean-popul. ages of 1.5-18 Gyr
- Coma E+S0 have mean-pop. ages 15-18 Gyr

4.
INTERACTIONS
AT HIGH REDSHIFT
AND IN CLUSTERS

Stockton + Farnham '91

Recent Merger in QSO OX 169 ($z=0.21$)




Results:


Tidal tail of
 $l_p \approx 30 \text{ kpc}$

($H_0 = 75$
 $q_0 = 0$)

$20'' \approx 60 \text{ kpc}$



Tail spectrum: Stellar




Variable $H\beta$ components
 $\rightarrow$ 2 different engines ! (?)

Radio Galaxy MRC 0406-244 ($z=2.44$)


Rush,
McCarthy,
etal 1997

Double
nucleus
Two central
light "tails"
(~ 30 kpc)

Two Ly α
bubbles



Mergers in Hercules Cluster ($z=0.034$)



(Las Campanas 2.5m plate, Alan Dressler)

CONCLUSIONS

- Gravitational interactions and mergers are shaping and reshaping galaxies throughout the observable universe. The vast majority involve **gas** and **dissipation**.
- The close link between mergers, ULIGs, and QSOs suggests that, like QSO activity, major merging may have **peaked around $z = 2$** .
- **Major disk-disk mergers** form **ellipticals** with kinematic subsystems and bimodal globular-cluster populations. Such mergers occurred relatively early in rich clusters and continue to the present day in poorer clusters and the field.
- **Minor mergers** tend to move disk galaxies toward earlier morphological types, creating kinematic subsystems and some **bulges**.
- **Hubble's morphological sequence appears to be mainly a sequence of decreasing merger damage.**

